



CL-Quant Add-on Module

A method for evaluating hPSC phenotype by measuring hPSC colony numbers using "hPSC Colony Count"

< BioStation CT usage example >

Benefits

- Using the CL-Quant image analysis software with the CL-Quant Add-on Module "hPSC Colony Count" makes it possible to count colonies of human pluripotent stem cells (hPSCs) during culturing.
- To monitor forming colony numbers during hPSCs culturing is important to confirm cell phenotypic characterization. Colony forming ratio is different also in different culture conditions. hPSCs are well known to change their phenotype with genetic change after prolonged culturing and usually increase their colony forming ratio. To monitor colony numbers at every passage during culture is important and useful to confirm their cell phenotype using CL-Quant Add-on Module "hPSC Colony count".

Observation device

- BioStation CT (Nikon, MLA10000)

Software

- CL-Quant ver. 5.02 (Nikon, MLS21000)

CL-Quant Add-on Module

- MA-PC-4X-CO01 hPSC Colony Count (Nikon, MLS30101)

Cells

- Human induced pluripotent stem (human iPS) cell line, Tic-FX (JCRB Cell Bank, JCRB1331.01, MRC-5-derived, Substrain of Tic iPS cell (JCRB1331) adapted to feeder-free culture)

Materials

- TeSR™-E8™ Kit for hESC/hiPSC Maintenance (STEMCELL Technologies, 05990)
- Fibronectin bovine plasma (Merck, F1141)
- StemPro™ EZPassage™ Disposable Stem Cell Passaging Tool (Thermo Fisher Scientific, 23181010)
- Cell Scraper S (Sumitomo Bakelite, MS-93100)
- Costar® 6-well Clear TC-treated Multiple Well Plates (Corning, 3516)

Methods

Human iPS cells, Tic-FX cells which were maintained in a medium of TeSR™-E8™ on a fibronectin-coated 25cm² flask without feeder cells were inoculated using Ezpassage™ and cell scraper. The cell clumps were seeded into the wells of a 6-well plate coated with Fibronectin and were cultured for 3 days in a BioStation CT at 37°C in a humidified atmosphere of 5% CO₂.

Phase contrast images of 8×8 fields (approx. 16.0 mm×16.0 mm) at the center of a well were automatically captured every 6 hours using a 4× objective lens, in a BioStation CT every 12 hours from 3 days after seeding. Auto-focusing was adjusted on the first field of view for a well and then images were automatically taken using that auto-focusing setting within that well.

The obtained image data were analyzed using CL-Quant and the Add-on Module "hPSC Colony Count" to automatically measure the colony count. The masked images of the area where human iPS cell colonies were localized were confirmed on the operation screen and output in Microsoft Excel® format.

Results

The analysis results using CL-Quant with the Add-on Module "hPSC Colony Count" display the colony area of Tic-FX cells cultured in TeSR™-E8™ in random coloring. Figure 1 shows an unprocessed phase contrast image of Tic-FX and its masked image after processing using CL-Quant with the Add-on Module "hPSC Colony Count". By displaying the areas recognized as colonies in random colors, it was possible to confirm the areas of individual iPSC colonies. The colony count was obtained automatically via CL-Quant and the Add-on Module.



Colony number	Phase contrast image	Masked image
64		

Figure 1. Phase contrast and masked images of hPSC colonies

Phase contrast image of Tic-FX human iPSC cell (left) was proceeded to mask colony areas with random colors (right). Colony number was obtained by automatic calculation in CL-Quant.

Summary

- Using the combination of CL-Quant and the Add-on Module "hPSC Colony Count" makes it possible to recognize individual areas of human iPSC colonies. The identified colonies are counted automatically and displayed on the operation screen.
- Monitoring the cell status during culturing hPSCs, and obtaining the information of the colony numbers can be used to assess the quality of hPSCs.
- The obtained numeric measured values can be output in Microsoft Excel® format.

< Introducing Nikon's observation systems >

The BioStation CT is equipped with an incubator for long-term monitoring of cells via microscope, and the BioStudio-T allows capturing without moving the stage. Both reduce stress on the cells and allow time-lapse photography of changes over time. Using Nikon's live cell imaging equipment and unique image analysis technology enables observation and real-time analysis of cell characteristics over time.



BioStation CT



BioStudio-T

ISO 14001 Certified
for NIKON CORPORATION



NIKON CORPORATION

Shinagawa Intercity Tower C, 2-15-3, Konan, Minato-ku, Tokyo 108-6290, Japan
phone: +81-3-6433-3705 fax: +81-3-6433-3785
www.healthcare.nikon.com/ja/

NIKON INSTRUMENTS INC.

1300 Walt Whitman Road, Melville, N.Y. 11747-3064, U.S.A.
phone: +1-631-547-8500: +1-800-52-NIKON (within the U.S.A. only)
fax: +1-631-547-0306
www.microscope.healthcare.nikon.com/ja_JP/